A situation in which the preimage of any maximal ideal stays maximal

written by rapha on Functor Network original link: https://functor.network/user/2593/entry/1113

For a general ring homomorphism, the preimage of a prime ideal stays prime. The situation is more complicated for maximal ideals: it's not true in general that a maximal ideal will stay maximal when pulled back! For a cheap example, look at the injection $\mathbb{Z} \to \mathbb{Q}$, for which the (maximal) ideal of \mathbb{Q} gets pulled back to the *minimal* ideal of \mathbb{Z} . This technical advantage of prime ideals over maximal ideals was one of the driving forces behind scheme theory.

In this post, we will explore one particular situation where maximal ideals are pullback-friendly: when the rings are k-algebras, and the target is of finite type.

The notation k will always denote some (any) field. Rings are commutative with unity.

* * *

Before we begin, we need a small technical lemma. Interesting in its own right, we will use this result to conclude that an integral domain is a field as soon as it admits a structure of a finite-dimensional vector space.

Lemma. If A is an integral domain that is also a finite algebra over k, then A is, in fact, a field.

Proof. Let a be some non-zero element of A. The main idea is to show that the function $\lambda_a: x \mapsto ax$ is a k-linear automorphism. To do this, it suffices to show that λ_a is an injection, because A is a finite-dimensional vector space. But this is clear, since A is an integral domain. \blacksquare

Parenthetically, notice that any integral domain admits a trivial vector space structure (multiplication by any scalar does nothing). Hence, if the integral domain has a finite number of elements, it is trivially a finite-dimensional vector space: by the lemma, it is a field!

Now we state the main point of this post:

Proposition. Let $\phi: A \to B$ be a morphism of algebras over k, and suppose further that B is of finite type over k. Then the preimage by ϕ of any maximal ideal of B is a maximal ideal of A.

In schematic terms, the induced morphism $\operatorname{Spec} B \to \operatorname{Spec} A$ maps closed points to closed points, so it corresponds more closely to one's intuition of what a morphism of varieties "should" be!

Proof. Suppose \mathfrak{m} is a maximal ideal of B. We want to show that the quotient ring $A/\phi^{-1}(\mathfrak{m})$ is a field. In general, the preimage of any maximal ideal is prime, so we already know $A/\phi^{-1}(\mathfrak{m})$ is an integral domain. We'll show that it is also a finite algebra over k, and then apply the lemma.

The morphism ϕ induces a canonical map $\widetilde{\phi}: A/\phi^{-1}(\mathfrak{m}) \to B/\mathfrak{m}$. It fits in the following commutative diagram of k-algebras and k-algebra homomorphisms:

$$\begin{array}{ccc} A & \xrightarrow{\hspace{0.5cm} \phi \hspace{0.5cm}} & B \\ \downarrow & & \downarrow \\ A/\phi^{-1}(\mathfrak{m}) & \xrightarrow{\hspace{0.5cm} \widetilde{\phi} \hspace{0.5cm}} & B/\mathfrak{m} \end{array}$$

Let's investigate the kernel of $\widetilde{\phi}$. Pick an arbitrary element \overline{v} in $A/\phi^{-1}(\mathfrak{m})$ (where $v \in A$ and the overline denotes passing to the quotient, as usual). We have

$$\overline{v} \in \ker \widetilde{\phi} \iff \widetilde{\phi}(\overline{v}) = 0 = \overline{\phi(v)}$$

$$\iff \phi(v) \in \mathfrak{m}$$

$$\iff v \in \phi^{-1}(\mathfrak{m})$$

$$\iff \overline{v} = 0.$$

Hence the kernel is trivial, that is, $\widetilde{\phi}$ is an injective morphism.

As a last step, recall Zariski's lemma (sometimes also called the Nullstellensatz, although it seems to me like this term is overloaded!): if a field K is of finite type as an algebra over some other field k, then K is in fact finite over k, i.e. $\dim_k K < \infty$. In our case, applying $K = B/\mathfrak{m}$ gives that B/\mathfrak{m} is actually a finite-dimensional vector space over k. But now $A/\phi^{-1}(\mathfrak{m})$ is isomorphic to a subspace of B/\mathfrak{m} via the injection $\widetilde{\phi}$, so it is also a finite-dimensional vector space over k.