

Abelian Galois representation implies potentially good reduction

[J'ignore](#)

10 Sep 2025

First we need a criterion for potentially good reduction in terms of Galois representation. The following is Theorem 2 from [Serre & Tate's paper *Good Reduction of Abelian Varieties*](#):

The abelian variety A has potential good reduction at a place v of K if and only if the image by $p : G_K \rightarrow T_\ell(A)$ of the inertia group $I(v)$ is finite.

Now the main result we aim to prove is the following (Corollary 1):

Suppose that the residue field k is finite of characteristic p , and that, for some $\ell \neq p$, the image of G_K in $\text{Aut}(T_\ell(A))$ is abelian. Then A has potential good reduction at v .

First we can complete K WLOG (since the Tate module of A is isomorphic to that of A' where A' is base change of A to the completion of K at v so if A' has potentially good reduction at v so is A by the above criterion). Local class field theory tells us that the image of inertia is actually a quotient of the group of units U_K of K . But structure theorem of unit groups of local fields tells us that U_K is the product of a finite group and a pro- p group. Thus the image intersect the pro- ℓ group $1 + \ell \cdot \text{End}(T_\ell)$ trivially. Hence the image inject into $\text{Aut}(T_\ell/\ell T_\ell)$, which is finite.