

# James reduced product construction

J'ignore • 13 Oct 2025

James reduced product  $J(X)$  is the free monoid generated by elements of  $X$  in the category of topological spaces. We would like to find out its homology. Note that  $J_i(X)$  is obtained from  $J_{i-1}(X)$  by the following **pushout diagram**.

$$\begin{array}{ccc}
 (X^{i-1})^{\vee i} & \longrightarrow & J_{i-1}(X) \\
 \downarrow & & \downarrow \\
 X^i & \longrightarrow & J_i(X) \\
 \downarrow & & \downarrow \\
 X^{\wedge i} & \longrightarrow & X^{\wedge i}
 \end{array}$$

Thus we see that  $J_i(X)$  is  $J_{i-1}(X)$  with  $X^{\wedge i}$  attached. To compute the homology of  $J_i(X)$  inductively, we note that if we take (reduced) suspension of this diagram (which remains a pushout since  $\Sigma$  being left adjoint preserves colimit), the vertical column splits since  $\Sigma(X \times Y) \cong \Sigma(X) \vee \Sigma(Y) \vee \Sigma(X \wedge Y)$ . Thus  $\Sigma J_i(X) \cong \Sigma J_{i-1}(X) \vee \Sigma(X^{\wedge i})$ . Thus we have

$$H_*(J(X)) \cong H_{*+1}(\Sigma J(X)) \cong H_{*+1}\left(\bigvee_i \Sigma(X^{\wedge i})\right) \cong \bigoplus_i H_*(X^{\wedge i}) \cong \bigoplus_i \tilde{H}_*(X)$$

The last isomorphism follows from Kunneth and  $H_*(X/A) \cong H_*(X, A)$  if  $A \rightarrow X$  is a cofibration (which implies that  $H_*(X \wedge X)$  is the quotient of  $H_*(X \times X)$  by  $H_*(X \times \{*\}) \oplus H_*(\{*\} \times X)$ ).

To compute the cohomology ring of  $J(S^{2k})$ , note that from its cellular structure we know that  $H^*(J(S^{2k}))$  has exactly one  $\mathbb{Z}$  at degree  $2ki$  for every  $i \geq 0$ . To figure out the ring structure, let  $e_{2ki} \in H^*(J(S^{2k}))$  be a generator of the degree  $2ki$  component of  $H^*(J(S^{2k}))$ . We first compute  $e_{2k}^i$  for any  $i \geq 0$ . Note that this is a multiple of  $e_{2ki}$ , say  $e_{2k}^i = c_i e_{2ki}$  for some  $c_i \in \mathbb{Z}$ . Let  $\pi : (S^{2k})^i \rightarrow J_i(S^{2k})$  be the attaching map, which induces  $\pi^* : H^*(J_i(S^{2k})) \rightarrow H^*((S^{2k})^i)$ . We will compute  $c_i$  via relating the image of  $e_{2k}^i$  and  $e_{2ki}$  under  $\pi^*$ . Note that by Kunneth the cohomology ring of  $H^*((S^{2k})^i)$  is  $E(a_{2k,j}, j = 1, \dots, i)$ , the exterior algebra with  $i$  generators at degree  $2k$ . It is easy to see that the image of  $e_{2ki}$  under  $\pi^*$  is  $\prod_{j=1}^i a_{2k,j}$ . To compute  $\pi^*(e_{2k}^i)$ , note that the preimage of  $S^{2k} \subset J_i(S^{2k})$  under  $\pi$  is  $(S^{2k})^{\vee i}$ , and it is easy to see from this that  $\pi^*(e_{2k}^i) = \sum_{j=1}^i a_{2k,j}$ . Thus

$$\pi^*(e_{2k}^i) = \left( \sum_{j=1}^i a_{2k,j} \right)^i = c_i \prod_{j=1}^i a_{2k,j}$$

Since  $a_{2k,j}^2 = 0$  and  $a_{2k,j}$  pairwise commute, we deduce that  $c_i = i!$ .

From knowing  $e_{2k}^i = i!e_{2ki}$  we show  $e_{2ki}e_{2kj} = \binom{i+j}{i}e_{2k(i+j)}$  easily. This implies the (integral) cohomology of  $J(S^{2k})$  is the divided power algebra on  $2k$ -dimensional class.

Reference: [https://en.wikipedia.org/wiki/James\\_reduced\\_product](https://en.wikipedia.org/wiki/James_reduced_product)