

# Equivariant derived categories

J'ignore · 9 Feb 2026

A principal  $G$ -bundle  $\pi : X \rightarrow Y$  need not be Zariski locally trivial, e.g. see [here](#), but it is at least etale-locally trivial. The idea is that every point  $y \in Y$  admits an etale neighborhood over which  $\pi$  has a right inverse.

A stronger claim is true if  $G$  is affine: Every principal bundle is locally isotrivial. This means the etale neighborhood can be chosen to be finite over its image in  $Y$  (i.e. the number of points in the fiber is constant, see [here](#)), e.g. see [this post](#).

An important proposition: Let  $P$  be a principal  $G$ -variety, and  $X$  another  $G$ -variety, then  $P \times X$  is a principal  $G$ -variety. For this we need  $X$  to be quasiprojective as well to ensure (\*) Every finite subset of  $X$  is contained in an affine open subset.

A  $G$ -equivariant sheaf on a  $G$ -variety  $X$  is a sheaf  $\mathcal{F}$  together with a rule  $\theta$  that assigns to every  $g \in G$  and  $U$  open an isomorphism  $\theta_{g,U} : \mathcal{F}(U) \rightarrow \mathcal{F}(g \cdot U)$ , and it should satisfy some associativity. Equivalently, we can say for every  $g \in G$ , there is a map  $\theta_g : \mathcal{F} \rightarrow \sigma_g^*(\mathcal{F})$  satisfying associativity where  $\sigma_g$  is the action of  $g$  on  $X$ . However, this doesn't take into account the topology of  $G$  (in other words we just treat  $G$  like a discrete group). A better definition is to view  $\theta_g$  as part of  $\theta : pr_2^* \mathcal{F} \rightarrow \sigma^* \mathcal{F}$  in  $Sh(G \times X, R)$ . Then the associativity axiom is  $b^* \theta \circ pr_{23}^* \theta = m^* \theta$  where  $m : G \times G \times X \rightarrow G \times X$  is multiplication of the first two coordinates,  $b : G \times G \times X \rightarrow G \times X$  is the action of the second coordinates, and  $pr_{23} : G \times G \times X \rightarrow G \times X$  is the projection onto the last two coordinates. The idea is that  $b^* \theta|_{\{g\} \times \{h\} \times X} = \sigma_h^* \theta_g$ ,  $\theta_h = pr_{23}^* \theta|_{\{g\} \times \{h\} \times X}$ , and  $\theta_{gh} = m^* \theta|_{\{g\} \times \{h\} \times X}$ , so this boils down to the usual axiom. The Proposition is that  $Sh_G(X, R)$  is an abelian category.

There are certain subtlety relating to equivariant derived category. It is neither  $D_c^b Perv_G$  nor just copy the above definition but allowing  $\mathcal{F}$  to be a general object in  $D_c^b$ . See section 6.4. We want the definition to satisfy the desideratum 6.1, including forgetful functor and six-functor formalism that is mutually compatible. See example 6.4.2 that uses the below proposition.

(Prop. 6.2.13) Every  $G$ -equivariant perverse sheaf on a  $G$ -homogeneous space  $X$  is a shifted local system and there is an equivalence of categories  $Loc_G^{ft}(X, k) \cong k[G_x / (G_x)^\circ] \text{-mod}^{fg}$ .

However, if  $X$  is a principal  $G$ -variety with geometric quotient  $\pi : X \rightarrow X/G$ , then we can define  $D_G^b(X, k)$  as  $D_c^b(X/G, k)$ . The key is Proposition 6.2.10 asserting  $\pi^\dagger$  induces an equivalence of categories between  $Perv(X/G)$  and  $Perv_G$ . This is just smooth descent for perverse sheaves plus identifying descent datum with the definition of  $G$ -equivariant sheaves using the Cartesian diagram for principal  $G$ -varieties.

For general  $G$ -varieties we use acyclic  $G$ -resolutions. See Exercise 3.7.2 for the definition of  $n$ -acyclic morphism. The key point is that for  $n$ -acyclic smooth morphism  $f : X \rightarrow Y$  the shifted pullback  $f^\dagger : {}^pD_c^b(Y, k)^{[k, k+n]} \rightarrow {}^pD_c^b(X, k)^{[k, k+n]}$  is fully faithful (easily seen from the adjunction). The motivation is that the cohomology of the fibers of  $f$  can lead to ‘extra’ morphisms in  $D_c^b(X, k)$  or in the category of descent data so there is no smooth descent on the level of derived category. However, if  $f$  is  $\infty$ -acyclic, then for all  $\mathcal{F} \in D_c^b(Y, k)$ , the unit and counit map  $\mathcal{F} \rightarrow f_*f^*\mathcal{F}$  and  $f_!f^!\mathcal{F}$  are both isomorphisms.

Lemma 6.1.20 tells us when  $X \rightarrow pt$  is an  $n$ -acyclic morphism in terms of  $H^i(X; \mathbb{Z})$ . The idea is to reduce to the case of constant sheaf  $\underline{\mathbb{Z}}$  on  $X$  and apply Exercise 3.2.1 (t-exactness of  $\otimes^L$  under good situations). From this we can produce lots of  $n$ -acyclic  $\mathbb{G}_m$  resolutions and  $GL_k$ -resolutions more generally. See Example 6.1.21 and 6.1.22. For linear algebraic group  $G$  this suffices to conclude the existence of  $n$ -acyclic  $G$ -resolution of any  $G$ -variety  $X$  using Prop. 6.1.13 and Prop. 6.1.15.

Now the definition for  $D_G^b(X, k)$  as in Definition 6.4.4 appears very natural. Essentially for every  $G$ -resolution  $P \rightarrow X$  we give an object  $\mathcal{F}(p) \in D_c^b(P/G, k)$  and it should be compatible for different  $G$ -resolutions in the sense that if  $\nu : P \rightarrow Q$  is a morphism of  $G$ -resolution then there is an isomorphism  $\alpha_\nu : \bar{\nu}^*\mathcal{F}(q) \rightarrow \mathcal{F}(p)$ . Moreover, the isomorphism  $\alpha_\nu$  should compose.

Note that we always have the trivial  $G$ -resolution  $pr_2 : G \times X \rightarrow X$  which allows us to define the forgetful functor  $\mathcal{F} \mapsto \mathcal{F}(pr_2)$ . It is not evidently a triangulated category with heart  $Perv_G$ . See [Berstein-Lunt’s book](#) for detailed proof.

Reference: [Achar’s book](#)