

The strong perfect graph theorem

Bogdan Grechuk · 17 Apr 2026

This post continues my series on favourite theorems of the twenty-first century. For an overview of the categories and earlier selections, see [this post](#).

My choice for 2002 in Combinatorics—and indeed across all of mathematics—is the celebrated strong perfect graph theorem of Chudnovsky, Robertson, Seymour, and Thomas. The theorem was proved in 2002 and published in 2006 (Chudnovsky et al. 2006).

The *chromatic number* of a graph G , denoted $\chi(G)$, is the smallest number of colours needed to colour the vertices of G so that adjacent vertices receive different colours. Such an assignment is called a *proper colouring*. Determining chromatic numbers for finite *simple* graphs—that is, graphs with no loops or multiple edges—is a deep and often difficult subject. A basic lower bound for $\chi(G)$ is the size $\omega(G)$ of the largest *clique* in G , since all vertices in a clique must receive distinct colours. In general, however, this bound need not be sharp. For example, if C is an odd cycle of length $n = 2k + 1 \geq 5$ (an *odd hole*), then $\omega(C) = 2$ while $\chi(C) = 3$. Likewise, if \bar{C} is the complement of such a cycle (an *odd antihole*), then $\omega(\bar{C}) = k$ but $\chi(\bar{C}) = k + 1$.

A graph G is called *perfect* if every induced subgraph H of G satisfies

$$\chi(H) = \omega(H).$$

A central problem in the area was to find a simple structural characterization of perfect graphs. The examples above show immediately that a perfect graph cannot contain an odd hole or an odd antihole as an induced subgraph. In 1961, Berge conjectured that this obvious necessary condition is also sufficient. This became known as the *strong perfect graph conjecture*. It attracted enormous attention and resisted proof for more than forty years.

Chudnovsky, Robertson, Seymour, and Thomas finally settled the conjecture in (Chudnovsky et al. 2006).

Theorem 1 *A graph G is perfect if and only if it does not contain an odd hole or an odd antihole as an induced subgraph.*

This is one of those rare classification theorems whose statement is both clean and unexpectedly definitive. Perfect graphs are defined by a condition involving the chromatic number of every induced subgraph, yet Theorem 1 shows that the

entire class can be recognized by forbidding just two families of induced subgraphs. The contrast between the complexity of graph colouring in general and the elegance of this criterion is, to me, part of what makes the theorem so beautiful.

The theorem also had major algorithmic consequences. Building on it, Chudnovsky, Cornuéjols, Liu, Seymour, and Vušković (Chudnovsky et al. 2005) developed a polynomial-time algorithm for recognizing perfect graphs. More precisely, they showed that there exists a polynomial P and an algorithm that, given a graph G on n vertices, performs at most $P(n)$ operations and determines whether G is perfect.

This fits perfectly with an earlier breakthrough of Grötschel, Lovász, and Schrijver (Grötschel et al. 1988), who gave a polynomial-time method for computing the chromatic number of a perfect graph. Taken together, these results mean that, for any graph G , we can determine in polynomial time whether G is perfect, and, if it is, compute $\chi(G)$ efficiently.

References

- Chudnovsky, Maria, Gérard Cornuéjols, Xinming Liu, Paul Seymour, and Kristina Vušković. 2005. “Recognizing Berge Graphs.” *Combinatorica* 25 (2): 143–86.
- Chudnovsky, Maria, Neil Robertson, Paul Seymour, and Robin Thomas. 2006. “The Strong Perfect Graph Theorem.” *Ann. of Math.* 164 (1): 51–229.
- Grötschel, Martin, László Lovász, and Alexander Schrijver. 1988. “Geometric Algorithms and Combinatorial Optimization.” *Algorithms Combin.* 2: 65–84.