

First post

User 3527 · 20 Oct 2026

<https://functor.network/user/3527/entry/2017>

Circle Tangency Criterion for Cyclic Quadrilateral

****Author:**** Aviraj Das ****Class:**** 9th Grade student, West Bengal, India

Abstract

We present a bidirectional characterization of cyclic quadrilaterals based on the tangency condition of two localized subcircles generated by diagonal intersections, orthocenter, and a circumcenter.

1. Introduction

The structural properties of cyclic quadrilaterals form a rich chapter in Euclidean geometry. This paper presents a necessary and sufficient condition for a convex quadrilateral to be cyclic: two constructed circles are tangent to each other at the diagonal intersection point.

2. Notations and Definitions

Let $ABCD$ be a convex quadrilateral whose diagonals AC and BD intersect at Q .

- O is the circumcenter of triangle QCD . - N is the midpoint of QC , and M is the midpoint of QD . - ω_1 is the circle through N, M, O, Q , with center α . - H is the orthocenter of triangle QAB . - G is the foot of the perpendicular from B onto QA . - F is the foot of the perpendicular from A onto QB . - ω_2 is the circle through H, Q, G, F , with center β . - I is the foot of the altitude from Q onto AB . - ω is the circumcircle of triangle QCD .

3. Main Theorem

****Theorem.**** The quadrilateral $ABCD$ is cyclic if and only if the circles ω_1 and ω_2 are tangent to each other at Q .

In for