

# Vector Sum and Minkowski Sum

written by Nada Ali on Functor Network


original link: <https://functor.network/user/409/entry/135>


**Theorem 1** (*Brunn – Minkowski inequality for convex bodies II*). *Let  $K$  and  $L$  be two convex bodies (i.e. compact convex sets with nonempty interiors) in  $\mathbb{R}^n$ , then:*

$$V(K + L)^{1/n} \geq V(K)^{1/n} + V(L)^{1/n} \quad (1)$$

In geometry, the term *Minkowski sum* is more frequently used for the vector sum. We now give an example to further clarify this claim.

Consider the origin-symmetric rectangle  $H$  of sides  $a$  and  $b$  and the other origin-symmetric rectangle  $S$  of sides  $c$  and  $d$  and their *Minkowski sum* shown below.





Now we compute their vector sum to show how it is equivalent to the *Minkowski sum*  $H + S$ :

$$V(H + S) = V(H) + V(S) + 2a\left(\frac{d}{2}\right) + 2b\left(\frac{c}{2}\right) \quad (2)$$

$$= V(H) + V(S) + ad + bc \quad (3)$$

$$\geq V(H) + 2\sqrt{V(H)V(S)} + V(S) \quad (4)$$

$$= (V(H)^{\frac{1}{2}} + V(S)^{\frac{1}{2}})^2 \quad (5)$$

So we get:

$$V(H + S)^{\frac{1}{2}} \geq V(H)^{\frac{1}{2}} + V(S)^{\frac{1}{2}} \quad (6)$$

which is *Minkowski* inequality (1) for  $n = 2$ .

*Remark.* Notice that we managed to derive equation (8) from equation (7) in the following way:

$$\begin{aligned} (ad + bc)^2 &\geq 0 \\ (ad)^2 + 2abcd + (bc)^2 &\geq 0 \\ a^2d^2 + b^2c^2 &\geq 2abcd \\ a^2d^2 + 2abcd + b^2c^2 &\geq 4abcd \\ \frac{(ad)^2 + 2abcd + (bc)^2}{4} &\geq abcd \\ \left(\frac{ad + bc}{2}\right)^2 &\geq V(H)V(S) \\ ad + bc &\geq 2\sqrt{V(H)V(S)} \end{aligned} \quad (7)$$